INDEX

Abattoirs, insect control about, 269–277.
Abutilon moth, 536.
Acanthoscelides obtectus, 190–193.
Acmaeodera angelica, 329.
  hepburnii, 329.
  mariposa, 330.
Aedes canadensis, 517, 518.
  sylvestris, 517, 518.
Agrilus acutipennis, 331.
  angelicus, 330.
  anxius, 331.
  granulatus, 331.
  niveiventris, 331.
  politus, 331, 332.
Ainslie, Geo. G., 114–123.
Aleurothrixus howardi, 377.
Aleurodicus minimus, 377.
Aleyrodes citri, 377.
Alfalfa weevil, 123–131.
Altitude and latitude law, 159–160, 161.
Alypia octomaculata, 47–48.
American Association Economic Entomologists, Proceedings, 1.
  Pacific Slope Branch, proceedings, 305–76.
  Secretary's report, 3–5.
Ammonia and fly, 532.
Amphiscepa bivittata, 552–556.
Anastatus (bifasciatus) 178, 179, 180, 181, 182.
Anemia, infectious of horses, 114.
Angoumois grain moth, 298.
Anopheles maculipennis, 517.
  occidentalis, 355.
  pseudopunctipennis, 355, 356.
  punctipennis, 517.
Anthaxia aenogaster, 328.
Anthonomus grandis, 312.
  signatus, 287–290.
Anthrenus scrophulariæ, 341
Apate francisca, 516.
Apeteticus maculiventris, 551.
Aphelinus lapisligni, 415.
Aphid eggs, 223–224, 556.
Aphidius testaceipes, 233, 237, 240.
Aphids and fire blight, 45–46.
Aphis avenæ, 556, 560.
  davisi, 418.
  gossypii, 313.
  neo-mexicana, var pacifica, 293.
  pomi, 556, 560.
Aphis rumexicolens, 417.
  saliceti, 417.
  sorbi, 556–560.
  viburniphila, 416.
Apiary inspection, 195–210.
Apple and thorn skeletonizer, 502.
Apple tree borer, round-headed, 66–71.
Arsenates, lead, 385–392.
Arsenic as an insecticide, 345–348.
Asphondylia websteri, 562.
Asterochiton abutiloneus, 313.
Atta texana, 561.
Attagenus plebius, 340–344.

Back, E. A., 453–458.
Baker, A. C., 420–433, 504.
Ball, E. D., 135–138.
Batrachedra rileyi, 445.
Bean weevil, 31, 32, 33, 37.
Becker, G. G., 49–59, 66–71.
Bee inspection, 200–203.
Bishopp, F. C., 269–277.
Biting lice, little red, 447.
Blaberus discoidalis, 224.
Blackhead fireworm, 553.
Blaps mucronata, 414.
Blatella germanica, 561.
Blister rust, white pine, 277–278.
Boncquet, P. A., and Stahl, C. F., 392–397.
Brassolis isthmia, 473–488.
Britton, W. E., 109–111.
Brown-tail moth, 193–195.
Bruchus chinensis, 74.
  obtectus, 31, 32, 190–193.
  quadrimaculatus, 74.
Buprestidæ, western, 325–332.
Buprestis confluens, 327.
Buffalo gnat, 413.
Burgess, A. F., and Griffin, E. L., 131–134.
Burke, H. E., 325–332, 406.

Caligonus mali, 499.
Callidium antennatum, 93.
Calliphora coloradensis, 273.
  erythrocephala, 273.
  iridescens, 273.
  vomitaria, 273.
Callipterinella annulata, 292, 427.
Camptocladius sp., 472–473.
Carbon dioxide and fly, 533.
  disulphide, 78.
Carr, E. G., 197–200.
Cattle flies, 111–113.
  lice, 446.
Ceratitis capitata, 318–321, 525.
Ceutorhynchus marginatus, 278–282.
Chaitophorus, key to species, 430.
  americanus, 428.
Chaitophorus bruneri, 429.
  lyropicta, 428.
  viminalis, 429.
Chalcis ovata, 542.
Chittenden, F. H., 282–287.
Chloridea virescens, 539.
Chlorochroa ligata, 309.

Chrysobothris femorata, 328.
mali, 328, 332.
Chrysocharis livida, 514.
Chrysomyia macellaria, 273.
Chrysophana placida, 330, 406.
Citrus mealy bug, 262.
Clerid larva and codling moth, 461–464.
Clover leaf weevil, 445.
Coccus citricola, 373–376.
hesperidum, 373–376.
Cockerell, T. D. A., 448, 562.
Cockroaches, 561.
Cocoanut-tree caterpillar, 473–488.
Codling moth, 60–63.
Cœlinidea meromyzæ, 527–531.
Cœliodes punctiger, 280.
Coffee insects, 513–517.
Coleman, G. A., 371–373.
Collins, C. W., 170–176.
Cooley, R. A., 94–102.
Cory, E. N., 111–113.
Cosmophila erosa, 536–542.
Cotton pests, 307–317.
Cottony cushion scale, 298.
Crambid moths, 114–123.
Crambus hortuellus, 553.
Cranberry girdler, 553.
rootworm, 553.
vinehopper, 552.
Crumb, S. E. and Lyon, S. C., 532–536.
Crepidodera rufipes, 414.
Crosby, C. R., and Leonard, M. D., 20–25.
Crossman, S. S., 177–183, 453–458.
Culex pipiens, 517.
restuans, 517.
tarsalis, 517.
Curly-top, beet, 392–397.
Cymatodera æthiops, 462, 463.
Cynomyia cadaverina, 273.
Cryptochætum monophlebi, 298.

Danaus archippus, 448.
Davidson W. M., 290–297, 350–353.
Davis, I. W., 193–195.
Davis, J. J., 41–44.
Dean, G. A., 146–159.
Dendroctonus monticolæ, 92.
de Ong, E. R., see Gray, Geo. P.
Dermacentor albipictus, 560.
Dermestes vulpinus, 340.
Dicerca hornii, 326.
Diplazon lætatorius, 529.
Diprion simile, 188–190, 224.
Dozier, H. L., 536–542.
Dunn, L. H., 473–488.
Dysdercus albidiventris, 309.

Eight spotted forester, 47–48.
Employment bureau, 6, 7.
Ephestia kuehniella, 446.
Epicranion championi, 516.
Epomphaloides minutus, 445.
Eriophyes thurberiæ, 314.
Essig, E. O., 433–444.
Euceraphis, key to species, 426.
betulæ, 425.
brevis, 426.
deducta, 427.
lineata, 426.
Euryopthalmus succinctus, 309.
Euschistus servus, 309.
Eutettex tenella, 392, 396.
Ewing, H. E., 497–501.

Fannia canicularis, 273.
Farm bureau and insect control, 20–25.
Felt, E. P., 60–63, 502, 562.
Ferris, G. H., 321–325.
Fire blight and aphids, 45–46.
Fly control exhibit, 411.
Foul brood, American, 414; European, 200.
Fox, Henry, 162.
Freeborn, S. B., 354–359.
Fruit flies, 333.
Fungus beetle, two-banded, 282–287.
growing ant, 561.

Galerucella luteola, 504.
Garman, H., 413.
Garman, Phillip, 503.
Gelechia gossypiella, 225, 315.
Gibson, E. H., 445, 503.
Gillette, C. P., 338–340.
Gipsy moth, 193–195; banding material, 131–134; wind dispersion, 170–176.
Glasgow, Hugh, 59.
Grain moth, 32.
Grasshopper control, 135–138; poison, 524–525.
Gray, G. P., 385–392, 353.
Gray, Geo. P., and de Ong, E. R., 353.
Green-bug, 233–248.

Hadley, C. H. and Matheson, R., 38–40.
Hadwen, Seymour, 447.
Hæmatobia serrata, 113.
Hæmatopinus eurysternus, 446.
vituli, 446.
Hawley, I. M., 545–552.
Hayes, W. P., 253–261.
Headlee, T. J., 31–38, 287–290.
Heliothrips fasciatus, 314.
Hemerocampa leucostigma, 175.
Hemerophila pariana, 502.
Hemiptera collection (U. S.) 502; workers directory, 446.
Hen lice, 71–74.
Herms, W. B., 359–70, 407–411, 563.
Hessian fly, 146–159, 162–168, 169, 249–253.
Hewitt, C. Gordon, 81–91.
Hop aphis, 551.
red bug, 545–552.
Horticultural board, 217–219.
inspection, 210–223.
Howard, C. W., 114, 411, 464–68, 517–521. 560, 561.

Housefly hibernation, 464–468; response, 102–109, 532.
Howard, L. O., 505.
Howardia biclavis, 516.
Humidity and insect metabolism, 31–38.
Hymenia perspectalis, 446.
Hymenoptera of Connecticut, 300.
Hypera punctata, 225, 445.
Hypoaspis armatus, 499.
Hypoderma lineatum, 447.
Hyslop, J. A., 278–282.

Illingworth, J. F., 340–344.
Index economic entomology, 9.
Indian meal moth, 563.
Insect behaviour, 81–91.
Insect collections, preservation, 445.
Insect photography, 25–30.
Insecticides, purer, 225–226.
Insects on nursery stock, 219–223.
Isosoma investigations, 139–146.
Isosoma tritici, 140, 141, 142, 143, 144, 145.
Inspection weaknesses, 216–217.

Janus abbreviatus, 224.
Jones, T. H., 561.

Kelly, E. O. G., 139, 233–248, 527–531.
Kentucky notes, 413.

Laboulbenia formicarum, 447.
Lachnopus sp., 515.
Lachnus rosæ, 418.
Lamson, G. H., 71–74, 447.
Lasius niger var. americana, 447.
Laurel psyllid, 439–444.
Lead arsenates, 385–392.
Leonard, M. D., 20–25.
Leptoglossus zonatus, 310.
Leucoptera coffeella, 514.
Ligyrus gibbosus, 253–261.
rugiceps, 162.
Lime as an insecticide, 74–78.
Lovett, A. L., 333–337, 345–348.
Lowry, Q. S., 47–48.
Loxostege sticticalis, 543–544.
Lucilia cæsar, 273.
Lygus pratensis, 310.
Lyon, S. C., 532–536.

Malacosoma pluvialis, 333.
Malaria-mosquito survey, 359–370.
Mallow caterpillar, 536–542.
Mansonia perturbans, 517.
Manter, J. A., 190–193.
Marchand, Werner, 469–472.
Marcovitch, Simon, 81.
Marmara elotella, 488–496.
Marsh, H. O., 543–544.
Matheson, R., 38–40.
McColloch, J. W., 162–168, 183–187.
McGregor, E. A., 504.
Mealy bugs, methods, 321–325.
Mediterranean flour moth, 446.
Mediterranean fruit-fly, 318–321, 521.
Megastigmus aculeatus, 448.
Melanophila gentilis, 327.
Melanoplus differentialis, 310.
Mercurial ointment, 71–74.
Meromyza americana, 528, 529, 530, 531.
Merrill, D. E., 461–464.
Merrill, J. H., 45–46.
Metcalf, Z. P., 74–78.
Midge infested potatoes, 472–473.
Miris dolobrata, 114.
Miscible oil *vs.* fish oil soaps, 453–458.
Mites, new economic, 497–501.
Monellia, key to species, 424.
Monieziella bipunctata, 501.
Morrill, A. W., 307–317.
Mosquito control, 517–521.
meeting (N. J.), 226–229.
work, anti-, 109–111.
Mosquitos, Californian, 362–363.
Motion pictures, 371–373.
Moznette, G. F., 344.
Myochrous longulus, 504.
Myrmelachista ambigua var. ramulorum, 515.
Myzocallis, key to species, 423.
alnifoliæ, 421.
californicus, 421.
fumipennellus, 422.
punctatellus, 420.
tiliæ, 421.
Myzus ribifolii, 294.
ribis, 338–340.

Newcomer, E. J., 445.
Nicotine sulphate, 333–337; tests, 459–461.

O'Kane, W. C., 78.
Okra caterpillar, 536–542.
Onion thrips, 521.
Ophyra ænescens, 273.
Ormenis pygmæa, 516.
Ornithodoros megnini, 407.
Osborn, Herbert, 114.
Oscinis variabilis, 414.
Ox louse, long-nosed, 446.
Ox louse, short-nosed, 446.

Paracalocoris haweleyi, 545–552.
Paraleptomastix abnormis, 262, 263, 264.
Parasite cages, 525–527.
Parasite colonizing, 177–183.
Paratrioza cockerelli, 434–439.
Parks, T. H., 249–253, 524–525.
Parrott, P. J., 79–81.
Patch, E. M., 416–420, 472–473.
Peach tree borer, 49–59.
Peairs, L. M., 507.
Pegomyia fusciceps, 397–406.
Pellett, F. C., 200–203.
Pemberton, C. E., and Williard, H. F., 525–527.
Peterson, Alvah, 556–560.
Petroleum insecticides, 353.

Phillips, E. F., 204–210.
Phillips, W. J., 139–146.
Phormia regina, 273.
Phorodon humuli, 551.
Photography, insect, 25–30.
Phyllocœlis atra, 554.
Polistes americanum, 542.
Phytomyza aquilegiæ, 224.
Pine blister rust, 213–215.
Pink boll worm, 225.
Pink cornworm, 445.
Piophila casei, 273.
Plant lice, western, 290–297.
Plodia interpunctilla, 563.
Plum aphis, reddish brown, 503.
Plum aphis, reddish-brown, 350–353.
Pœcilonota cyanipes, 327.
ferrea, 327.
thureura, 327.
Polycesta californica, 329.
Prociphilus approximatus, 418.
xylostei, 418.
Pseudococcus citri, 262, 266.
longispinus, 516.
Psychonoctua jamaicensis, 516.
Pterocomma, key to species, 431.

Quarantine, importance of, 298.
Quayle, H. J., 373–376.

Radish maggot, 79–81.
Red-legged flea beetle, 414.
Reduviolus subcoleoptratus, 551.
Reeves, G. I., 123–131.
Rhabdopterus picipes, 553.
Rhinoncus pyrrhopus, 278.
Rhopalosiphum nymphææ, 350–353, 503.
Rhopobota vacciniana, 553.
Rice fields and malaria, 354–359.
Richardson, C. H., 102–109.
Rockwood, L. P., 415.
Root maggot, 397–406.

Safro, V. I., 459–461, 521–523.
Saissetia hemispherica, 516.
Sanders, J. G., 213–215.
Sanderson, E. D., 507.
Sanninoidea exitiosa, 49–59.
Sarcophaga cimbicis, 260.
helicis, 260.
rudis, 260.
Sasscer, E. R., 79, 219–223.
Scammell, H. B., 552–556.
Schedius kuvanæ, 178, 179, 180, 181, 182.
Schistocerca shoshone, 310,
vega, 310.
Schizotetranychus latitarsus, 498.
Schœne, W. J., 216–217.
School entomology, 507.
Seventeen-year locust, 38–40.
Severin, H. P., 318–321, 333.
Shaw, H. B., 217–219.
Silpha bituberosa, 94–102.
Simulium pecuarum, 413.
venustum, 413.
Sinuate pear borer, 59.
Sitotroga cerealella, 32, 298.
Smith, H. S., 262–268.
Smith, R. H., 447.
Spinach carrion beetle, 94–102.
Spinose ear tick, 407–411.
Spraying truck crops, 521–523.
Stahl, C. F., 392–397.
Stomoxys calcitrans, 113.
Strawberry weevil, 81, 287, 290.
Sugar-beet leafhopper, 392.
webworm, 543–544.
Sunflower insects, 561.

Tabanid rearing, 469–472.
Tarsenomus pallidus, 344, 501, 503.
Tetranychus bimaculatus, 314.
multidigituli, 497.
uniunguis, 497.
Tetrastichus xanthomelænæ, 504.
Tettigonia occatoria, 516.
Tibicen septendecim, 38–40.
Toadbugs, 554.
Toxoptera aurantii, 516.
graminum, 233–248.
outbreak, 139.
Tomato psyllid, 434–439.
Turner, W. B., 445.
Trachykele blondeli, 326.
Trichodectes scalaris, 447.
Trichogramma pretiosa, 542.
Trioza alacris, 439–444.
Tucker, E. S., 397–406.

Underground insects, 183–187.

Vacuum fumigation, 79.
Viereck, H. L., 300.
Vinal, S. C., 488–496.

Walden, B. H., 25–30.
Washburn, F. L., 277–278.
Wasmannia auropunctata, 515.
Wheat fly, 414.
White grubs, analysis of, 41–44.
Weiss, H. B., 224, 448.
Webster, R. L., 225.
Weevils, cow-pea, 74.
Williard, H. F., 525.
Wilson, T. S., 445.
Wind dispersion, gipsy moth, 170–176.

Yingling, H. C., 223–224.

Zagrammosoma multilineata, 514.
Zappe, M. P., 188–190.
Zwaluwenburg, R. H. Van, 513–517.